

Cetrilin™ - A

LevoCetirizine Dihydrochloride+ Pseudoephedrine+ Ambroxol

CETRILIN™ - ACetirizine Dihydrochloride + Pseudoephedrine
+ Ambroxol Tablets

PHARMA BUSINESS



WORLDWIDE

Cetrilin-A

Comsition

LevoCetirizine Dihydrochloride.....	5 mg
Pseudoephedrine.....	30 mg
Ambroxol.....	60 mg

Levocetirizine is a third-generation non-sedative antihistamine, developed from the second-generation antihistamine cetirizine. Chemically, levocetirizine is the active enantiomer of cetirizine. It is the L-enantiomer of the cetirizine racemate. Levocetirizine works by blocking histamine receptors. It does not prevent the actual release of histamine from mast cells, but prevents it binding to its receptors. This in turn prevents the release of other allergy chemicals and increased blood supply to the area, and provides relief from the typical symptoms of heigh fever.

Indications

Cetrilin-A is indicated in the treatment of perennial and seasonal allergic rhinitis pruritus, allergic asthma, allergic conjunctivitis and urticaria.

Description

An antihistamine is a drug serves to reduce or eliminate effects. which serves to reduce or eliminate effects mediated by histamine. an endogenous chemical mediator released during allergic reactions. Only agents where the main therapeutic effect is mediated by negative modulation of histamine receptors are termed antihistamines - other agents may have antihistaminergic action but are not true antihistamines. In common use, the term antihistamine refers only to H1-receptor antagonists, also known as H1-antihistamines.

Dosage and Administration

Adults and Children 12 yr of age and older

2.5 to 5 mg once daily in the evening (max, 5 mg once daily).

Children 6 to 11 yr of age

2.5 mg (½ tablet or 1 teason oral solution) once daily in the evening (max, 2.5 mg once daily).

Dose Adjustment for Renal Function Impairment

Adults and Children 12 yr of age and older

Mild renal function impairment

Recommended dosage is 2.5 mg once daily.

Moderate renal function impairment

Recommended dosage is 2.5 mg once every other day.

Severe renal function impairment

Recommended dosage is 2.5 mg twice weekly, separated by 3 to 4 days.



Taj Pharmaceuticals Limited, Taj Group Pharmaceuticals Division

Andheri (w), Maharashtra 4000 53, INDIA

Revised Every Year

6794553d

www.tajpharmaceuticals.com

www.tajagroproducts.com

www.tajfordoctors.com

Cetrilin™ - A

LevoCetirizine Dihydrochloride+ Pseudoephedrine+ Ambroxol

CETRILIN™ - ACetirizine Dihydrochloride + Pseudoephedrine
+ Ambroxol Tablets

PHARMA BUSINESS



WORLDWIDE

Cetrilin-A

Indications and Usage

Relief of symptoms associated with allergic rhinitis; treatment of uncomplicated skin manifestations of chronic idiopathic urticaria.

Contraindications

Patients with end-stage renal disease and patients undergoing hemodialysis; children 6 to 11 yr of age with renal function impairment; hypersensitivity to any component of the product or to cetirizine.

Presentations

10 tablets

Note : This product information is intended only for residents of the India. Taj Pharmaceuticals Limited, medicines help to treat and prevent a range of conditions—from the most common to the most challenging—for people around the world.

Information for Health Care Professionals

*** Please consult local Prescribing Information for any product before use. This website is an international information resource for healthcare professionals with an interest in disease management. This website is not intended to replace the advice of a qualified healthcare professional. Above brand is a trademark of the Taj group of companies (Taj Pharmaceuticals Limited).



Taj Pharmaceuticals Limited, **Taj Group Pharmaceuticals Division**

Andheri (w), Maharashtra 4000 53, INDIA

Revised Every Year

6794553d

www.tajpharmaceuticals.com

www.tajagroproducts.com

www.tajfordoctors.com