




Betasec™

Betahistine dihydrochloride
8/16/24/32 MG

Rx Vertigo



Vertigo is usually associated with a problem in the inner ear balance mechanisms (vestibular system), in the syndrome and Meniere's disease.

Transient **vertigo can be caused by inflammation of the inner ear due to the common cold, influenza, and bacterial infections. Vertigo can also occur after long flights or boat journeys where the mind gets used to turbulence, resulting in a person's feeling as if he or she is moving up and down.** This usually subsides after a few days. Another source of vertigo is through exposure to high levels of sound pressure, rattling the inner ear and causing a loss of balance. Consumption of alcohol can also cause vertigo.

Rarely, vertigo-like symptoms may appear as **paraneoplastic syndrome (PNS)** in the form of opsoclonus myoclonus syndrome, an extremely rare multi-faceted neurological disorder associated with many forms of incipient cancer lesions or viruses.

- **Available as Tablets**
- Uses : Vertigo
- **Side effects*** Gastric upset, headache, rash, dizziness, sleeplessness

- Use with Care Bronchial asthma
- Do not give anti-histamines concurrently
- Not used in Pregnancy and breast feeding
- Children below 12 years
- Peptic ulcer



NOTE: * The side effects are seen in a small percentage of persons using the medicine. The benefits of using the drug usually out-weigh the side-effects. Your doctor is the best person to decide the use and dosage of any drug.

Betasec™
Betahistine dihydrochloride
8/16/24/32 MG