

Dilpril®
Enalapril Maleate

Tablet®
DILPRIL
Enalapril Maleate



Dilpril

USES

This drug belongs to a group of medications called ACE inhibitors. Dilpril® is used to treat high blood pressure (hypertension) in adults and children. It works by relaxing blood vessels, causing them to widen. Lowering high blood pressure helps prevent strokes, heart attacks and kidney problems. This medication is also used with other drugs (e.g., "water pills"/diuretics, digoxin) to treat congestive heart failure.

DOSAGE

Take this medication by mouth, usually once or twice a day; or as directed by your doctor. You may take this drug with or without food. Use this medication regularly in order to get the most benefit from it. Remember to use it at the same time(s) each day.

If you are taking this drug in the liquid suspension form, shake the bottle well before each use. Measure the dose out carefully.

Do not take potassium supplements or salt substitutes containing potassium without talking to your doctor or pharmacist first. This medicine can raise your potassium levels, which rarely can cause serious side effects such as muscle weakness or a very slow heartbeat. Tell your doctor immediately if either of these effects occur.

The dosage is based on your medical condition and response to therapy. The usual maximum dose for adults is 40 milligrams per day.

Dilpril® side effects

- *feeling light-headed, fainting;
- *urinating more or less than usual, or not at all;
- *fever, chills, body aches, flu symptoms;
- *pale skin, easy bruising or bleeding;
- *fast, pounding, or uneven heartbeats;
- *chest pain; or
- *swelling, rapid weight gain.



Dilpril®
Tablet


DILPRIL Tablet®
Enalapril Maleate



Dilpril

Less serious Dilpril ® side effects

- *cough;
- *loss of taste sensation, loss of appetite;
- *dizziness, drowsiness, headache;
- *sleep problems (insomnia);
- *dry mouth;
- *nausea, vomiting, diarrhea; or
- *mild skin itching or rash.

STORAGE

Store tablets at room temperature below 86 degrees F (30 degrees C) away from light and moisture. Do not store in the bathroom. Refrigerate the liquid suspension between 36-46 degrees F (2-8 degrees C).

Note : This product information is intended only for residents of the India. Taj Pharmaceuticals Limited, medicines help to treat and prevent a range of conditions—from the most common to the most challenging—for people around the world.

Information for Health Care Professionals

*** Please consult local Prescribing Information for any product before use. This website is an international information resource for healthcare professionals with an interest in disease management. This website is not intended to replace the advice of a qualified healthcare professional. Above brand is a trademark of the Taj group of companies (Taj Pharmaceuticals Limited).



Taj Pharmaceuticals Limited, **Taj Group Pharmaceuticals Division**

Andheri (w), Maharashtra 4000 53, INDIA

Revised Every Year

6794553d

www.tajpharmaceuticals.com

www.tajagroproducts.com

www.tajfordoctors.com