

Fetrasil™
Sibutramine hydrochloride monohydrate

FETRASIL™
(Sibutramine Hydrochloride Monohydrate)



Fetrasil

Each Fetrasil Tablets contains 5 mg, 10 mg, and 15 mg of sibutramine hydrochloride monohydrate. It also contains as inactive ingredients: lactose monohydrate, NF; microcrystalline cellulose, NF; colloidal silicon dioxide, NF; and magnesium stearate, NF in a hard-gelatin Tablets [which contains titanium dioxide, USP; gelatin; FD&C Blue No. 2 (5- and 10-mg Tabletss only)

Fetrasil is indicated for the management of obesity, including weight loss and maintenance of weight loss, and should be used in conjunction with a reduced calorie diet. Fetrasil is recommended for obese patients with an initial body mass index = 30 kg/m², or = 27 kg/m² in the presence of other risk factors (e.g., diabetes, dyslipidemia, controlled hypertension).

DOSAGE AND ADMINISTRATION

The recommended starting dose of Fetrasil is 10 mg administered once daily with or without food. If there is inadequate weight loss, the dose may be titrated after four weeks to a total of 15 mg once daily. The 5 mg dose should be reserved for patients who do not tolerate the 10 mg dose. Blood pressure and heart rate changes should be taken into account when making decisions regarding dose titration (see WARNINGS and PRECAUTIONS).

Doses above 15 mg daily are not recommended. In most of the clinical trials, Fetrasil was given in the morning.

Storage

Store at 25°C (77°F); excursions permitted to 15°-30°C (59°-86°F) [see USP controlled room temperature]. Protect Tabletss from heat and moisture. Dispense in a tight, light-resistant container as defined in USP.

WARNINGS

Blood Pressure and Pulse

Fetrasil SUBSTANTIALLY INCREASES BLOOD PRESSURE AND/OR PULSE RATE IN SOME PATIENTS. REGULAR MONITORING OF BLOOD PRESSURE AND PULSE RATE IS REQUIRED WHEN PRESCRIBING Fetrasil.



Fetrasil™
Sibutramine hydrochloride monohydrate

FETRASIL™
(Sibutramine Hydrochloride Monohydrate)



Fetrasil

PRECAUTIONS

Pulmonary Hypertension

Certain centrally-acting weight loss agents that cause release of serotonin from nerve terminals have been associated with pulmonary hypertension (PPH), a rare but lethal disease. In premarketing clinical studies, no cases of PPH have been reported with sibutramine Tablets. Because of the low incidence of this disease in the underlying population, however, it is not known whether or not Fetrasil may cause this disease.

Seizures

During premarketing testing, seizures were reported in < 0.1% of sibutramine treated patients. Fetrasil should be used cautiously in patients with a history of seizures. It should be discontinued in any patient who develops seizures.

Bleeding

There have been reports of bleeding in patients taking sibutramine. While a causal relationship is unclear, caution is advised in patients predisposed to bleeding events and those taking concomitant medications known to affect hemostasis or platelet function.

Presentation

Loravan Tablets Blister of 10 Tablets

Note : This product information is intended only for residents of the India. Taj Pharmaceuticals Limited, medicines help to treat and prevent a range of conditions—from the most common to the most challenging—for people around the world.

