



## Hito-EN®

### HITO-EN Syrup

Cyproheptadine Hydrochloride (anhydrous) IP .... 2 mg

#### INDICATIONS:

HITO-EN Syrup are indicated in the treatment of acute and chronic allergies such as dermatitis, eczema, mild and local allergic reactions to insect bites, hay fever, seasonal rhinitis, allergic conjunctivitis due to inhalant allergens and foods, urticaria, angioneurotic oedema, drug and serum reactions, anogenital pruritus, and pruritus of chicken-pox.

HITO-EN Syrup may also be used in the treatment and prophylaxis of migraine and vascular headaches.

#### DOSAGE AND ADMINISTRATION

##### Allergy and Pruritus

###### Adults

12–16 mg per day in divided doses. The dosage should not exceed 32 mg daily.

###### Children (Aged 7–14 Years)

Usually 4 mg thrice daily. The dosage should not exceed 16 mg daily.

###### Children (Aged 2–6 Years)

2 mg thrice daily. Total dosage should not exceed 12 mg daily.

###### Children below 2 Years of Age

Not recommended

##### Migraine and Vascular Headaches

Prophylactically or therapeutically, the recommended dosage is 4 mg initially, repeated after half-hour if necessary, but not to exceed 8 mg in a 4–6 hour period. In responsive patients, relief is usually obtained with two doses (totally 8 mg) and maintained with 4 mg every 4-6 hours.





## Hito-EN®

### CONTRAINDICATIONS

Angle-closure glaucoma, stenosing peptic ulcer or pyloroduodenal obstruction, symptomatic prostatic hypertrophy or bladder-neck obstruction, concurrent monoamine oxidase inhibitor therapy, acute asthmatic attacks, and hypersensitivity to cyproheptadine.

### PACKAGING INFORMATION

HITO-EN Syrup .....Bottle of 100 ml

Presentations: 100ml

