

Metherlin®
(methylergonovine maleate)
Tablets, USP

Metherlin®
(methylergonovine
maleate) Tablets, USP
0.2 mg



TAJ PHARMACEUTICALS LIMITED

“Working for healthier India”

DRUG DESCRIPTION

Metherlin (methylergonovine maleate) is a semi-synthetic ergot alkaloid used for the prevention and control of postpartum hemorrhage.

Active Ingredient: methylergonovine maleate, USP, 0.2 mg.

Inactive Ingredients: acacia, carnauba wax, D&C Red #7, FD&C Blue #1, gelatin special, lactose, maleic acid, mixed parabens, povidone, sodium benzoate, sodium hydroxide, starch, stearic acid, sucrose, talc, and titanium dioxide.

INDICATIONS

For routine management after delivery of the placenta; postpartum atony and hemorrhage; subinvolution. Under full obstetric supervision, it may be given in the second stage of labor following delivery of the anterior shoulder.

USES

This medication is used to help stop bleeding after delivery of the placenta in childbirth. Methylergonovine maleate belongs to a class of drugs known as ergot alkaloids. It works by increasing the stiffness of the uterus muscles after the last stage of labor. This effect decreases bleeding.

DOSAGE AND ADMINISTRATION

Intramuscularly

0.2 mg, after delivery of the anterior shoulder, after delivery of the placenta, or during the puerperium. May be repeated as required, at intervals of 2-4 hours.

Intravenously

Dosage same as intramuscular.

Orally

One tablet, 0.2 mg, 3 or 4 times daily in the puerperium for a maximum of 1 week.

R

Taj Pharmaceuticals Limited, Taj Group Pharmaceuticals Division
Andheri (w), Maharashtra 4000 53, INDIA
Revised Every Year
6794553d



tajpharmaceuticals.com
tajagroproducts.com
www.tajfordoctors.com

Metherlin[®]
(methylergonovine maleate)
Tablets, USP

Metherlin[®]
(methylergonovine
maleate) Tablets, USP
0.2 mg



TAJ PHARMACEUTICALS LIMITED

“Working for healthier India”

SIDE EFFECTS

The most common adverse reaction is hypertension associated in several cases with seizure and/or headache. Hypotension has also been reported. Nausea and vomiting have occurred occasionally. Rarely observed reactions have included: acute myocardial infarction, transient chest pains, arterial spasm (coronary and peripheral), bradycardia, tachycardia, dyspnea, hematuria, thrombophlebitis, water intoxication, hallucinations, leg cramps, dizziness, tinnitus, nasal congestion, diarrhea, diaphoresis, palpitation, rash, and foul taste.¹

There have been rare isolated reports of anaphylaxis, without a proven causal relationship to the drug product.

Drug Abuse And Dependence

Metherlin (methylergonovine maleate) has not been associated with drug abuse or dependence of either a physical or psychological nature.

WARNINGS

This drug should not be administered I.V. routinely because of the possibility of inducing sudden hypertensive and cerebrovascular accidents. If I.V. administration is considered essential as a lifesaving measure, Metherlin (methylergonovine maleate) should be given slowly over a period of no less than 60 seconds with careful monitoring of blood pressure. Intra-arterial or periarterial injection should be strictly avoided.

PRECAUTIONS

Before taking methylergonovine maleate, tell your doctor or pharmacist if you are allergic to it; or to similar ergot alkaloids (e.g., ergonovine); or if you have any other allergies.

This medication should not be used if you have certain medical conditions. Before using this medicine, consult your doctor or pharmacist if you have: high blood pressure, a certain complication of pregnancy (toxemia).

Before using this medication, tell your doctor or pharmacist your medical history, especially of: heart disease (e.g., venoatrial shunt, mitral valve stenosis), other complications during pregnancy (e.g., pre-eclampsia, eclampsia), a serious blood infection (sepsis), blood vessel problems (e.g., Raynaud's phenomenon, obliterative vascular disease), kidney problems, liver problems.

This drug may make you dizzy or drowsy. Use caution while driving, using machinery, or doing any activity that requires alertness. Limit or avoid alcoholic beverages.

R

Taj Pharmaceuticals Limited, **Taj Group Pharmaceuticals Division**
Andheri (w), Maharashtra 4000 53, INDIA
Revised Every Year
6794553d



tajpharmaceuticals.com
tajagroproducts.com
www.tajfordoctors.com

Metherlin®
(methylergonovine maleate)
Tablets, USP

Metherlin®
(methylergonovine
maleate) Tablets, USP
0.2 mg



TAJ PHARMACEUTICALS LIMITED

“Working for healthier India”

STORAGE

Store at room temperature below 77 degrees F (25 degrees C) away from light and moisture. Do not store in the bathroom. Keep all medicines away from children

Note : This product information is intended only for residents of the India. Taj Pharmaceuticals Limited, medicines help to treat and prevent a range of conditions—from the most common to the most challenging—for people around the world.

Information for Health Care Professionals

*** Please consult local Prescribing Information for any product before use. This website is an international information resource for healthcare professionals with an interest in disease management. This website is not intended to replace the advice of a qualified healthcare professional. Above brand is a trademark of the Taj group of companies (Taj Pharmaceuticals Limited).

R

Taj Pharmaceuticals Limited, **Taj Group Pharmaceuticals Division**
Andheri (w), Maharashtra 4000 53, INDIA
Revised Every Year
6794553d



tajpharmaceuticals.com
tajagroproducts.com
www.tajfordoctors.com