



Amlace

Each 5 mg Tablet Contains

Composition

Amlodipine5 mg

INDICATIONS

Hypertension

AMLACE® is indicated for the treatment of hypertension. AMLACE is indicated for the symptomatic treatment of chronic stable angina. AMLACE may be used alone or in combination with other antianginal agents.

AMLACE® is indicated for the treatment of confirmed or suspected vasospastic angina. AMLACE® may be used as monotherapy or in combination with other antianginal drugs.

In patients with recently documented CAD by angiography and without heart failure or an ejection fraction <40%, AMLACE is indicated to reduce the risk of hospitalization due to angina and to reduce the risk of a coronary revascularization procedure.

Use

Amlodipine (Amlace®) is in a group of drugs called calcium channel blockers. It works by relaxing (dilating) your blood vessels, lowering blood pressure, and decreasing heart rate, which lowers the workload on the heart. It also dilates coronary arteries increasing blood flow to the heart.

Amlodipine is used to treat high blood pressure (hypertension) or chest pain (angina).

Dosage

Take Amlace® exactly as prescribed by your doctor. Do not take it in larger amounts or for longer than recommended. Follow the directions on your prescription label.

Amlace® is usually taken once daily. Your doctor may occasionally change your dose to make sure you get the best results from this medication. Your chest pain may become worse when you first start taking amlodipine or when your dose is increased. Call your doctor if your chest pain is severe or ongoing.

If you are being treated for high blood pressure, keep using this medication even if you feel fine. High blood pressure often has no symptoms, so you may not know when your blood pressure is high. You may need to use blood pressure medication for the rest of your life.

Store amlodipine at room temperature away from moisture, heat, and light.





Amlace

Side Effects

Amlodipine may cause side effects. Tell your doctor if any of these symptoms are severe or do not go away:

- * swelling of the hands, feet, ankles, or lower legs
- * headache
- * upset stomach
- * stomach pain
- * dizziness or lightheadedness
- * drowsiness
- * excessive tiredness
- * flushing (feeling of warmth)

Some side effects can be serious. The following symptoms are uncommon, but if you experience any of them, call your doctor immediately:

- * more frequent or more severe chest pain
- * rapid, pounding, or irregular heartbeat
- * fainting

Presentation

Amlace® 10 Tablet per Blister

Note : This product information is intended only for residents of the **India**. Taj Pharmaceuticals Limited, medicines help to treat and prevent a range of conditions—from the most common to the most challenging—for people around the world.

Information for Health Care Professionals

*** Please consult local Prescribing Information for any product before use. This website is an international information resource for healthcare professionals with an interest in disease management. This website is not intended to replace the advice of a qualified healthcare professional. Above brand is a trademark of the Taj group of companies (Taj Pharmaceuticals Limited).

