



Combi Fast Plus

Composition

Paracetamol IP, Phenylephine, Triprolidine

Properties

Drug combined with analgesic, antipyretic, decongestant and expectorant action.

Paracetamol inhibits cyclooxygenaze and reduces prostaglandins synthesis, pain and temperature. Ascorbic acid compensates vitamine C loss that is characteristic for the incipient phases of acute respiratoty infections.

Phenylefrin contracts sanguine vessels and reduces congestion.

Terpinhydrate stimulates bronchial secretion and facilitates expectoration. Caffeine increases the analgesic effect of paracetamol.

Prescribed for

diseases provoked by cold, influenza.

Contraindications

Hyper sensibility to any of the drug components, severe renal and hepatic affections, arterial hypertension, hyperthyrosis, diabetes, heart diseases.

Dosing

Adults - 2 tablets (maximum 4 tablets) per day. 6-12 year old children - 1 tablet 4 times per day at the most.

Side effects

Allergic reaction (cutaneous eruption, angioneurotic edem, rash). Rarely arterial hyper tension (cephalalgia, nausea, vomiting, diarrhea, insomnia, palpitations).

Drug interactions

The drug is incompatibile with tricyclic antidepressant (risk of hypertensive thurst) and betablockants.

Cautions

pregnancy and lactation period, children younger than 3 years.