

**Gynameno-Plus®**  
Tranexamine Acid , Mefenamic Acid.

# Gynameno-Plus ®



## Gynameno-Plus

**Each tablet contains**

### Composition

Tranexamine Acid .....500 mg  
Mefenamic Acid.....250 mg

### Indications

For powerful control of painful bleeding.

### Description

Tranexamic acid (commonly marketed as Cyclokapron) is often prescribed for excessive bleeding. It is an antifibrinolytic that competitively inhibits the activation of plasminogen to plasmin. Tranexamic acid is commonly used to treat excessive menstrual bleeding, even to the point of menorrhagia. It is also useful in the treatment of haemophilia.

Mefenamic acid binds the prostaglandin synthetase receptors COX-1 and COX-2, inhibiting the action of prostaglandin synthetase. As these receptors have a role as a major mediator of inflammation and/or a role for prostanoid signaling in activity-dependent plasticity, the symptoms of pain are temporarily reduced.

### Contraindications

Mefenamic acid: is contraindicated in patients with active ulceration or chronic inflammation of either the upper or lower gastrointestinal tract. It should not be used in patients with preexisting renal disease.

### Dosage

As per the physician's advice.

### Presentations

10 tablets

**Note :** This product information is intended only for residents of the India. Taj Pharmaceuticals Limited, medicines help to treat and prevent a range of conditions—from the most common to the most challenging—for people around the world.

### Information for Health Care Professionals

\*\*\* Please consult local Prescribing Information for any product before use. This website is an international information resource for healthcare professionals with an interest in disease management. This website is not intended to replace the advice of a qualified healthcare professional. Above brand is a trademark of the Taj group of companies (Taj Pharmaceuticals Limited).



# Gatimin ®



## Gatimin

Acute pyelonephritis  
400 mg once daily for 7-10 days

Gonorrhoea in women.  
400 mg as a single dose

### **Contraindications**

Gatimin is contraindicated in persons with a history of hypersensitivity to gatifloxacin or any member of the quinolone class of antimicrobial agents.

### **Dosage**

Dosages of Gatimin are administered once every 24 hours.

### **Presentations**

10 tablets

**Note :** This product information is intended only for residents of the India. Taj Pharmaceuticals Limited, medicines help to treat and prevent a range of conditions—from the most common to the most challenging—for people around the world.

### **Information for Health Care Professionals**

\*\*\* Please consult local Prescribing Information for any product before use. This website is an international information resource for healthcare professionals with an interest in disease management. This website is not intended to replace the advice of a qualified healthcare professional. Above brand is a trademark of the Taj group of companies (Taj Pharmaceuticals Limited).

