



TAJ PHARMACEUTICALS LIMITED

"Working for healthier India"

It is a broad spectrum anticonvulsant

Mechanism of Action

Valproate appears to act by multiple mechanisms:

- (a) A Phenytoin like frequency dependent prolongation of Na⁺ channel inactivation
- (b) Attenuation of Ca²⁺ mediated 'T' current
- (c) Augmentation of release of inhibitory transmitter GABA by inhibiting its degradation as well as increasing its synthesis.

Oral absorption of Valproic acid is good. Peak plasma levels are achieved in 1-4 hrs. It is 90% bound to plasma proteins; 95% metabolized in liver by oxidation and glucuronide conjugation – excreted in urine. Plasma t_{1/2} is 15 hours; but anticonvulsant effects are longer lasting

Indications

Valproic acid is highly effective in absence seizures and is an alternative / adjuvant drug for generalized and partial seizures

To prevent recurrent febrile seizures in children, also used in some cases of Myoclonic and Atonic seizures

Dosage

Adults : 200mg 3 times a day, gradually increased at weekly intervals upto a maximum of 600mg 3 times a day according to response.

Children : 10-15mg/kg/day in three divided doses, increase according to response at weekly interval by 5mg/kg/day upto 30-40mg/kg/day

Contraindications

Known hypersensitivity

Special Precautions

Caution in patients in hepatic and renal diseases
Monitor liver function test





TAJ PHARMACEUTICALS LIMITED

"Working for healthier India"

Adverse Drug Reactions

Anorexia, nausea, vomiting, drowsiness, ataxia are dose related effects Alopecia, hyperammonemia, sedation, vertigo, tremor, nystagmus and confusion can occur with prolonged use Hypersensitivity reactions like rashes and thrombocytopenia

Asymptomatic increase in serum transaminases, rarely hepatitis and pancreatitis It is teratogenic

Status in

1. Pregnancy: Contraindicated
2. Lactation : Use with caution
3. Old age : May be given in reduced dose
4. Children : May be given in dose as advised

Interactions

Phenytoin: Increases its level by inhibiting metabolism and displacing it from protein binding
Phenobarbital, primidone: Inhibits their metabolism
Clonazepam: Simultaneous administration may precipitate absence syndrome
Carbamazepine: Induce each other's metabolism

Presentation

Wallbrurate _____ Blister of 10 Tablets

Note : This product information is intended only for residents of the India. Taj Pharmaceuticals Limited, medicines help to treat and prevent a range of conditions—from the most common to the most challenging—for people around the world.

Information for Health Care Professionals

*** Please consult local Prescribing Information for any product before use. This website is an international information resource for healthcare professionals with an interest in disease management. This website is not intended to replace the advice of a qualified healthcare professional. Above brand is a trademark of the Taj group of companies (Taj Pharmaceuticals Limited).

