



Vasodin

Each Tablet contains:

Trimetazidine dihydrochloride tablet..... 35mg

COMPOSITION :

Boxes of 10 and 60 modified release film-coated tablets dosed at 35 mg of trimetazidine dihydrochloride. Excipients. q.s.f. one tablet.

THERAPEUTIC INDICATIONS

This drug is recommended in:

- * Preventive treatment for episodes of angina pectoris (Chronic stable angina).
- * Adjuvant symptomatic treatment for vertigo and tinnitus (noises detected by the ear whistling, buzzing, which do not correspond to any exterior sounds).
- * Adjuvant treatment of visual disorders of a circulatory origin.

CONTRAINDICATIONS :

Do not take Vasodin if you are allergic to any of the constituents.
This drug is generally not recommended during breast feeding

SPECIAL WARNINGS AND PRECAUTIONS :

This drug is not a curative treatment for angina attacks, nor an initial treatment for unstable angina pectoris. It is not a treatment for myocardial infarction. In the event of an angina attack, inform your doctor. Tests may be required and your treatment may possibly be modified.

SIDE EFFECTS :

As with all drugs, Vasodin is likely to have undesirable effects :

- * Rare cases of gastro-intestinal disorders (nausea, vomiting)
- * It you notice any undesirable effects, which have not been mentioned here, you should inform your doctor or pharmacist.





Vasodin

PREGNANCY AND BREASTFEEDING :

Pregnancy

It is preferable not to take this medicine during pregnancy, if you discover that you are pregnant whilst taking this medicine, consult your doctor as he alone can judge the necessity of continuing your treatment

Breastfeeding

In the absence of data on excretion in breast milk, breastfeeding is not recommended during treatment. Ask your doctor or pharmacist's advice before taking any medicine.

DOSAGE AND METHOD OF ADMINISTRATION

Oral route.

The usual dosage is 1 tablet in the morning and evening. Tablets should be swallowed with a glass of water at meal times.

If you think that the effect of Vasodin is too strong or too weak, talk to your doctor or pharmacist.

OVERDOSAGE :

Consult a doctor or a pharmacist immediately.

WHAT TO DO IN CASE ONE OR SEVERAL DOSES HAVE BEEN MISSED

Resume treatment normally. Do not take a double dose to compensate for the single dose that you forgot to take.

STORAGE CONDITIONS :

Keep out of the reach and sight of children.

Do not use after the expiry date indicated on the box.

